

BIBLIOGRAFÍA

Aberg JA, Gallant JE, Jean Anderson J, et al. Primary Care Guidelines for the Management of Persons Infected with Human Immunodeficiency Virus: Recommendations of the HIV Medicine Association of the Infectious Diseases Society of America. *CID* 2004; 39: 609-29. Disponible en www.journals.uchicago.edu/CID/journal

Accidente con material biocontaminado: esquema de actuación. Unidad de Salud Laboral, Servicio de Medicina Preventiva. Hospital Universitario Virgen Macarena.

Almeda J, Casabona J, Allepuz A, Garcia-Alcaide F, del Romero J, Tural C et al. Recomendaciones para la profilaxis postexposición no ocupacional al VIH.

American Academy of Pediatrics Committee on Pediatric AIDS; American Academy of Pediatrics Infectious Diseases and Immunization Committee. Evaluation and treatment of the human immunodeficiency virus-1-exposed infant. *Pediatrics* 2004;114:497-505

American Academy of Pediatrics, Committee on Nutrition. Cholesterol in childhood. *Pediatrics* 1998 ;101(1 pt 1):141-7.

ASRM. Hepatitis and reproduction. The Practice Committee of the American Society for Reproductive Medicine. *Fertil Steril* 2004;82:1754-1764.

Barreiro P, Duerr A, Beckerman K, Soriano V. Reproductive options in HIV-serodiscordant couples. *AIDS Rev.* 2006b;8:158-70.

Barreiro P, Castilla JA, Labarga P, Soriano V. Is natural conception a valid option for HIV-serodiscordant couples? *Hum Reprod* 2007;22:2353-58

Barreiro P, Del Romero J, Leal M, Hernando V, Asencio R, De Mendoza C, Labarga P, González-Lahoz J, Soriano V, Natural pregnancies in HIV-serodiscordant couples under successful antiretroviral therapy. *JAIDS* 2006a;43:324-6.

Bartlett JG and Gallant J. Medical management of HIV infection: 2005-2006. Baltimore, Maryland. The Johns Hopkins University, Division of Infectious Diseases; 2005.

Bartlett JG. Adult HIV/AIDS Treatment 2006. Disponible en www.hopkins-aids.org

Bascuñana Quirell A, Pueyo Rodríguez I. En: Pachón J, Rivero A, Pujol E, eds. La infección por el VIH. Guía práctica. Sevilla, 2002. Enfermedades de transmisión sexual e infección por el VIH: 313-322.

Benson CA, Kaplan JE, Masur H, Pau A, Holmes KK, CDC, National Institutes of Health, Infectious Diseases Society of America. Treating opportunistic infections among HIV-Exposed and Infected Children: Recommendations from CDC, the National Institutes of Health, and the Infectious Diseases Society of America. *Clin Infect Dis* 2005;40:S131-S235.

Bergshoeff A, Burger D, Farrelly L, et al. Pharmacokinetics of once daily vs twice daily lamivudine and abacavir in HIV-1 infected children: PENTA 13.

BHIVA guidelines writing Committee. Guidelines for the management of HIV infection in pregnant women and the prevention of mother-to-child transmission of HIV. *HIV Medicine* 2005;6:107-48. Disponible en: www.bhiva.org

British Columbia Health Service. Clinical Management of Chronic Hepatitis C. 2004. http://www.healthservices.gov.bc.ca/msp/protoguides/gps/hepatitis_c.pdf

British HIV Association and Children's HIV Association guidelines for the management of HIV infection in pregnant women 2008. *HIV Medicine* (2008), 9, 452–502

Bujan L, Hollander L, et al. For the CREAThE network. Safety and efficacy of sperm washing in HIV-1-serodiscordant couples where the male is infected: results from the European CREAThE network. *AIDS* 2007;21:1909-1914

Bujan L, Pasquier C, Labeyrie E, Lanusse-Crousse P, Morucci M, Daudin M. Insemination with isolated and virologically tested spermatozoa is a safe way for human immunodeficiency type 1 virus-serodiscordant couples with an infected male partner to have a child. *Fertil.Steril.* 2004;82:857-62.

Cabrera A, López JM, Blasco E, Salinas J. Guía práctica de manejo de exposiciones ocupacionales a VIH, VHB, VHC en base a las recomendaciones de los CDC. *Medicina Preventiva*, vol 3, 2, 2º trimestre, 2002; 26-30.

Centers for Disease Control and Prevention (CDC), Health Resources and Services Administration; National Institutes of Health, HIV Medicine Association of the IDSA. Incorporating HIV prevention into the medical care of persons living with HIV. Recommendations of CDC, the Health Resources and Services Administration, the National Institutes of Health, and the HIV Medicine Association of the IDSA. *MMWR Recomm Rep* 2003; 52(RR-12):1–24. Disponible en www.cdc.gov/mmwr

Centers for Disease Control and Prevention (CDC). 1994 revised classification system for HIV in children under 13 years of age. *MMWR* 1994;43:1-10.

Centers for Disease Control and Prevention. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HIV and Recommendations for Postexposure Prophylaxis. *MMWR* 2005;54(No. RR-9):1-17.

Centres for Diseases Control. Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV, and HIV and Recommendations for Postexposure Prophylaxis. *MMWR* 2001; 50 (No. RR-11).

Cervia JS. Transitioning HIV-infected children to adult care. *J Pediatr* 2007; 150: e1.

Chadwick E, Rodman J, Palumbo P, et al. A prospective evaluation of pharmacologic, virologic and immunologic parameters of lopinavir/ritonavir for HIV-1 infected infants below 6 months of age. 12th CROI Boston 2005. Abstract 766.

Comisión Asesora sobre Técnicas de Reproducción Humana Asistida del Departamento de Sanidad y Seguridad Social de la Generalitat de Cataluña. Técnicas de Reproducción Asistida y VIH. (CATRHAGC, 2002).

Committee on Infectious Diseases, The American Academy of Pediatrics. Pertussis. In: Pickering LK, ed. 2006 Red Book: Report of the Committee on Infectious Diseases. 27th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2006.

Dankner WM, Lindsey JC, Levin MJ; Pediatric AIDS Clinical Trials Group Protocol Teams 051, 128, 138, 144, 152, 179, 190, 220, 240, 245, 254, 300 and 327. Correlates of opportunistic infections in children infected with the human immunodeficiency virus managed before highly active antiretroviral therapy. *Pediatr Infect Dis J* 2001;20:40-48.

de Arístegui J, Corretger JM, Fortuny C, Gatell JM, Mensa J. Guía de Terapéutica Antimicrobiana en Pediatría 2004. Collbató (España) Ed. Antares; 2004.

Diabetes mellitus y Atención Primaria. Grupo de Diabetes de la Sociedad Andaluza de Medicina Familiar y Comunitaria, disponible en: [Document from the European Project on "Standardization of the management of occupational exposure to HIV/blood-borne infections and evaluation of post-exposure prophylaxis in Europe." FUNDING: European Commission, Directorate-General Health Care and Consumer Protection. Unit F4. Project number SI2.322294 Recommendations for Post-Exposure Prophylaxis against HIV infection in Health Care Workers in Europe. March 2002.](http://www.cica.es/aliens/samfyc/Brotons C, Royo-Bordonada MA, Álvarez-Sala L, Armario A, Artigao R, Conthe P, et al. Comité Español Interdisciplinario para la Prevención Cardiovascular (CEIPC). Adaptación española de la Guía Europea de Prevención Cardiovascular. Clin Invest Arterioscl 2005;17:19-33.</p></div><div data-bbox=)

Dore GJ, Kaldor JM. Detection of HCV RNA in semen. *Lancet* 2000;356:1520

Dornadula G, Zhang H, VanUitert B, Stern J, Livornese L, Ingerman MJ et al. Residual HIV-1 RNA in blood plasma of patients taking suppressive highly active antiretroviral therapy. *JAMA* 1999;282:1627-1632.

Dubé MP, Stein JH, Aberg JA, Fichtenbaum CJ, Gerber JH, Tashima KT, et al, for the Adult AIDS Clinical Trials Group Cardiovascular Subcommittee. Guidelines for the evaluation and management of dyslipidemia in human immunodeficiency virus (HIV)-infected adults receiving antiretroviral therapy: recommendations of the HIV Medicine Association of the Infectious Disease Society of America and the Adult AIDS Clinical trials Group. *Clin Infect Dis* 2003;37:613-627.

Duffaut M, Valla D. Procréation médicalement assistée et infection par le virus de l'hépatite C. *Contracept Fertil Sex* 1997;25:534-537

Dunn D. HIV Pediatric Prognostic Markers Collaborative Study Group. Short term risk of disease progression in HIV-1 infected children receiving no antiretroviral therapy or zidovudine monotherapy; a meta-analysis. *Lancet* 2003;362; 1605-11.

Educación para el paciente:

http://www.aidsinfo.nih.gov/ed_resources/default.php?cat_id=P

<http://www.medscape.com/patientdirectory/hiv?src=hdr>

<http://www.projectinform.org>

Englert Y, Lesage B, Van Vooren JP, Liesnard C et al. Medically assisted reproduction in the presence of chronic viral diseases. *Hum Reprod Update* 2004;10:149-162

European Collaborative Study. Mother-to-child Transmission of the era of Highly Active Antiretroviral Therapy. *Clin Infect Dis*. 2005;40: 458-65.

<http://www.aidsinfo.nih.gov/ContentFiles/PerinatalGL.pdf>

<http://www.prosego.com> Infección por VIH y Gestación

<http://www.rcog.org.uk/index.asp?PageID=8> Guidelines

European Paediatric Lipodystrophy Group. Antiretroviral therapy, fat redistribution and hyperlipidaemia in HIV-infected children in Europe. *AIDS* 2004;18:1443-51.

Expert panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive Summary of the third report of the National Cholesterol Education Program (NCEP)

Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *JAMA* 2001;285:2486-2497.

Faye A, Le Chenadec J, Dolffus C, et al. Early versus deferred antiretroviral multidrug therapy in infants infected with HIV type 1. *Clin Infect Dis*. 2004; 39:1692-99.

Fiscus SA, Brambilla D, Coombs RW et al., Multicenter evaluation of methods to quantitate human immunodeficiency virus type 1 RNA in seminal Plasma. *J Clin Micro* 2000;38:2348-2353.

Foster C, Mackie N, Seery P, Walters S, Tudor-Williams G, Lyall H. Emerging multi-drug resistance in children with perinatally acquired HIV-1. 8th International Congress on Drug Therapy in HIV Infection. Glasgow 2006. Abstract P360.

Galán I, Jiménez JL, González-Rivera M, De José MI, Navarro ML, Ramos JT, et al. Virological phenotype switch under salvage therapy with lopinavir-ritonavir in heavily pretreated HIV-1 vertically infected children: an additional benefit. *AIDS* 2004; 18: 247-55.

Gatell JM, Clotet B, Podzamczar B, et al. "Guía práctica del SIDA. Clínica, diagnóstico y tratamiento. 8ª edición". Ed Masson; 2004.

Giaquinto C, Notheis G, Della Negra M, Jelaska A, Miki J on behalf of the BI1182 study group. Efficacy of tipranavir oral solution co-administered with low dose ritonavir is similar in all age groups of children and teenagers after adjustment for resistance. 8th International Congress on Drug Therapy in HIV Infection. Glasgow 2006. Abstract P316.

Gilling-Smith C, Emiliani S, Almeida P, Liesnard C, Englert Y. laboratory safety during assisted reproduction in patients with blood-borne viruses. *Hum Reprod* 2005;20:1433-1438 .

Gona P, Van Dyke RB, Williams PL, Dankner WM, Chernoff MC, Nachman SA, Seage GR 3rd. Incidence of Opportunistic and Other Infections in HIV-Infected Children in the HAART Era. *JAMA* 2006;296:292-300

Gray R, Wayer M, Briikmeyer R, Sewankambo N et al. Probability of HIV-1 transmission per coital act in monogamous, heterosexual, HIV-1 serodiscordant couples in Rakai, Uganda. *Lancet* 2001 ;357:1149-1153.

Grupo de consenso español sobre profilaxis postexposición no ocupacional al VIH. *Enferm Infecc Microbiol Clin* 2002; 20(8):391-400.

Hawkins D, Blott M, Clayden P, de Ruiter A, Foster G, Gilling-Smith C, et al. BHIVA. Guidelines Writing Committee. Guidelines for the management of HIV infection in pregnant women and the prevention of mother-to-child transmission of HIV. *HIV Med.* 2005;6(Suppl 2): 107-48.

HIV InSite Database of Antiretroviral Drug Interactions.
<http://www.hivinsite.com/InSite.jsp?page=ar-00-02>

HIV Tools Drug Interaction Tool. <http://www.hivtools.com/>

http://www.sefh.es/01informacion_farmacoterapeutica_vih.php#1

Informe anual 2007. Registro andaluz de casos de SIDA. Monografía 2008. Sistema de Vigilancia Epidemiológica de Andalucía, vol. 13, nº17.

InteraccionesHIV.com. <http://www.interaccioneshiv.com/Buscar.php>

J. González-García et al. Coinfección por VIH y virus de las hepatitis A, B y C en pacientes adultos. Revisión y recomendaciones de GESIDA/PNS. *Enferm Infecc Microbiol Clin*; Febrero 2003 Disponible en www.gesidaseimc.com

Klein J, Pena JE, Thornton MH, Saber MV. Understanding the motivations, concerns and desires of human immunodeficiency virus 1-serodiscordant couples wishing to have children through assisted reproduction. *Obstet Gynecol* 2003;101:987-994.

La infección por el VIH: Guía Práctica. 2ª Edición. Capítulo 43. Profilaxis postexposición. Montserrat Pérez Pérez, Manuel Zarzuela Ramírez y Daniel Peña Jiménez

Labarga P, Martínez E, Soriano V, Barreiro P. Panel Español de Estudio de Parejas VIH-discordantes. Consejo reproductivo en parejas serodiscordantes para el virus de la inmunodeficiencia humana. *Med Clin (Barc)*. 2007;129:140-48

Lameiro FJ, Repáraz F, Sola J, Tiberio G, Pavón A, Gost J. Control de la infección en el personal sanitario I: inmunización. Enfermedades vehiculadas por sangre y secreciones. *Anales del Sistema Sanitario de Navarra*, suplemento 2.

Leonard EG, et al. Metabolic complications of antiretroviral therapy in children. *Pediatr Infect Dis J* 2003;22:77-84.

Lereuz-Ville M, Kunstmann JM, De Almeida M, Rouzioux C, Chaix ML. Detection of hepatitis C virus in the semen of infected men. *Lancet* 2000;356:42-43.

Liverpool Pharmacology Group. www.hiv-druginteractions.org

Lundgren JD, Battegay M, Behrens G, De Wit S, Guaraldi G, Katlama C, et al. European AIDS Clinical Society (EACS) guidelines on the prevention and management of metabolic diseases in HIV. *HIV Medicine* 2008;9:72-81.

Maertens A, Bourlet T, Plotton N, Pozzetto B, Levy R. Validation of safety procedures for the cryopreservation of semen contaminated with hepatitis C virus in assisted reproductive technology. *Hum Reprod*. 2004 Jul;19(7):1554-7.

Marina S, Marina F, Alcolea R et al. Human immunodeficiency virus type 1-serodiscordant couples can bear healthy children after undergoing intrauterine insemination. *Fertil Steril.* 1998;70:35-9.

Masur H, Kaplan JE, Holmes KK, U.S. Public Health Service, Infectious Diseases Society of America. Guidelines for preventing opportunistic infections among HIV-infected persons-2002. Recommendations of the U.S. Public Health Service and the Infectious Diseases Society of America. *Ann Intern Med* 2002;137:435-478.

Medscape Drug Interaction Calculator. <http://www.medscape.com/px/hivscheduler>

Meseguer, M, Garrido N, Gimeno C, Remohí J, Simón C, Pellicer A. Comparison of polymerase chain reaction: dependent methods for determining the presence of human immunodeficiency virus and hepatitis C virus in washed semen. *Fertil Steril* 2002; 78: 1199-1202.

MMWR. General Recommendations on Immunization. Feb,2002 / 51(RR02); 1-36 Immunizations for infected adults: indications, timing and response. *Immunizations*, vol 14, issue 5, dec 2006-jan 2007.

MMWR. Recommendations of the Advisory Committee on Immunization Practices (ACIP): Use of vaccines and Immune Globulins in Persons with altered immunocompetence. April, 1993, vol.42, No. RR-4

Monografías para pacientes. http://www.hivmedicationguide.com/pamphlets_Spanish.htm

Mora S, et al. Longitudinal changes of bone mineral density and metabolism in antiretroviral-treated human immunodeficiency virus-infected children. *J Clin Endocrinol Metab* 2004; 89: 24-28.

Mortimer D. Current and future concepts and practices in human sperm cryobanking. *RBMonline* 2004;9:134-151.

Nachman S, Gona P, Dankner W, Weinberg A, Yogev R, Gershon A, Rathore M, Read JS, Huang S, Elgie C, Hudgens K, Hughes W. The rate of serious bacterial infections among HIV-infected children with immune reconstitution who have discontinued opportunistic infection prophylaxis. *Pediatrics* 2005;115:e488-494.

Narayana K, D'Souza UJ, Rao S. Ribavirin-induced sperm shape abnormalities in Wistar rat. *Mutat Res* 2002;15:193-196.

OMS-ONUSIDA Antiretroviral therapy and sexual transmission of HIV. 1 febrero, 2008.

Pachón J, Pujol E, Rivero A. "La Infección por el VIH. Guía práctica. 2ª edición" Sevilla: Sociedad Andaluza de Enfermedades Infecciosas/Consejería de Salud; 2003. Disponible en www.saei.org

Página de cáncer de cérvix de los Centers for Disease Control and Prevention. <http://www.cdc.gov/cancer/cervical>

Página de infecciones de transmisión sexual de los Centers for Disease Control and Prevention. <http://www.cdc.gov/std/default.htm>

Página de la Sociedad Americana de Colposcopia y patología Cervical. <http://www.asccp.org>

Pérez-Milan F. Infección por virus de la Hepatitis C y Reproducción asistida. Boletín SEF 2002;10 (<http://nuevo.sefertilidad.com/boletin/2002/vol2/06.php>)

Perinatal HIV Guidelines Working Group. Recommendations for use of antiretroviral drugs in pregnant HIV-1-infected women for maternal health and interventions to reduce perinatal HIV-1 transmission in the United States. October 12, 2006. Disponible en <http://AIDSinfo.nih.gov>

Podzamczar D, Miralles P, de la Calle M, et al. Recomendaciones de GESIDA/Plan Nacional sobre el sida respecto al diagnóstico y tratamiento del sarcoma de Kaposi y cáncer de cérvix uterino en pacientes infectados por el virus de la inmunodeficiencia humana. Med Clin 2002;118:788-795.

Polo R, Galindo MJ, Martínez E, Alvarez J, Arévalo JM, Asensi V, et al; Study Group for Metabolic Alterations/Secretariat for the Nacional AIDS Plan (GEAM/SPNS). Recommendation of the Study Group for Metabolic Alterations/Secretariat for the Nacional AIDS Plan (GEAM/SPNS) of the management of metabolic and morphologic alterations in patients with HIV infection. Enferm Infecc Microbiol Clin 2006;24:96-117.

Procedimiento 004 del Sistema de Prevención de Riesgos Laborales del SAS.

Profilaxis postexposición parenteral frente al VIH y los virus B y C de la hepatitis. Servicio de Cuidados Críticos y Urgencias, Servicio de Enfermedades Infecciosas, Servicio de Farmacia Hospitalaria, Servicio de Medicina Preventiva y Salud Pública y Servicio de Microbiología. Hospitales Universitarios "Virgen del Rocío". Sevilla, Agosto 2001.

Public Health Service Task Force Recommendations for Use of Antiretroviral Drugs in Pregnant HIV-Infected Women for Maternal Health and Interventions to Reduce Perinatal HIV Transmission in the United States - July 8, 2008

Puthanakit T, Oberdorfer A, Akarathum N, et al. Efficacy of highly active antiretroviral therapy in HIV-infected children participating in Thailand's National Access to Antiretroviral Program. Clin Infect Dis 2005;41:100-7.

Quinn TC, Waver MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. N Engl J Med 2000;342:921-929.

Ramos JT, De José MI, Dueñas J, et al. Safety and efficacy of lopinavir/ritonavir in HIV-infected children experienced with 3 classes of antiretrovirals. Pediatr Infect Dis J 2005; 24: 867-73.

Recomendaciones de Gesida/Plan Nacional sobre el SIDA. http://www.gesida.seimc.org/pcientifica/fuentes/DcyRc/DcyRc_Recomendaciones_TAR_GESIDA-PNS_Impver1.pdf

Recomendaciones de GESIDA/Plan Nacional sobre el SIDA respecto al tratamiento antiretroviral en adultos infectados por el virus de la inmunodeficiencia humana (actualización enero de 2007). Panel de expertos de GESIDA y Plan Nacional sobre el SIDA. Enferm Infecc Microbiol Clín;25(1): 32-53. Disponible en www.gesidaseimc.com

Recomendaciones de la OMS: "Adherence to long- term therapies": Chapter XII: HIV and AIDS. http://www.who.int/chronic_conditions/adherencereport/en

Recomendaciones de la Secretaría del Plan Nacional sobre el SIDA (SPNS), el Grupo de Estudios sobre SIDA (GESIDA), la Sociedad Española de Ginecología y Obstetricia (SEGO) y la Asociación Española de Pediatría (AEP) para el seguimiento de la infección por el VIH con relación a la reproducción, el embarazo y la prevención de la transmisión vertical. 2007. Disponible en: www.msc.es

Recomendaciones para mejorar la adherencia de la SEFH. Documento de consenso.
<http://www.msc.es/profesional/preProSalud/sida/pdfs/adherencia.pdf>

Recomendaciones sobre reconocimientos médicos específicos de los trabajadores sanitarios con riesgo de exposición a virus de transmisión sanguínea. Ministerio de Sanidad y Consumo, 2004.

Recommended Adult Immunization Schedule, Oct 06-sep 07, CDC.
www.cdc.gov/nip/recs/adult-schedule.pdf

Sauer M, Choi J. HIV-seroconversion in a woman preparing for assisted reproduction: an inherent risk in caring for HIV-serodiscordant couples RBM on line 2006;12: 375-377

Sharland M, Blanche S, Castelli G, Ramos JT, Gibb D, on behalf of the PENTA steering committee. PENTA (Pediatric European Network for Treatment of AIDS). European guidelines of antiretroviral treatment of HIV-infected children. HIV Medicine 2004.

Sharland M, Castelli G, Ramos JT, Wintegest U, Gibb D, et al, on behalf of the PENTA steering committee. PENTA. European guidelines of antiretroviral treatment of HIV-infected children. HIV Medicine 2004 Foster C, et al. Emerging multi-drug resistance in children with perinatally acquired HIV-1. 8th International Congress on Drug Therapy in HIV Infection. Glasgow 2006. Abs. p360.

Sirera G, Videla S, Herranz P, Corzo-Delgado J, por el Grupo de Estudio del SIDA (GeSIDA). Virus del papiloma humano y virus de la inmunodeficiencia humana/SIDA. Enferm Infecc Microbiol Clin 2006;24(Supl.2): 40-47.

Sociedad Española de Medicina Familiar y Comunitaria. Hepatitis C. Recomendaciones. 2003

Speck CE, Coombs RW, Koutsky LA, Zeh J, Ross SO et al. Risk factors for HIV-I shedding in semen. Am J Epidemiol 1999;150:622-631

Standards of medical care in diabetes. Diabetes Care 2005;28: (Suppl 1): S4-36.

Steyaert SR, Leroux-Roels GG, Dhont M. Infections in IVF: review and guidelines. Hum Reprod Update 2000;6:432-441.

Sutinen J. Interventions for managing antiretroviral therapy-associated lipoatrophy. Curr Opin Infect Dis 2005;25-33.

Travelers' Health:: Yellow book. Health Information for International travel, 2005-2006. Vacunas recomendadas en adultos portadores de VIH. Manual de vacunaciones del adulto, 2005. Guide canadien d'immunisation, 6ème édition, 2002.

V Puro, S Cicalini, G De Carli, F Soldani, G Ippolito, on behalf of the European Occupational Post-Exposure Prophylaxis Study Group*. Towards a standard HIV post exposure prophylaxis for healthcare workers in Europe. Eurosurveillance monthly 2004; 9 (6):3-4.

Vacunación en adulto. Recomendaciones. Comisión de Salud Pública del 18 de Febrero de 2009.

van Leeuwen E, Prins JM, Jurriaans S, Boer K, Reijs P, Repping, van der Veen F. Reproduction and fertility in human immunodeficiency virus type-1 infection. *Hum Reprod Update* 2007;13:197-206

van Leeuwen E, Visser M, Prins JM, Nieuwkerk PT, van der Veen F. HIV couples' anxiety and risk taking during ART . *Fertil Steril* 2007a (epub ahead of print)

Vernazza P, Hirschel Bb, Bernasconi E, Flepp M. Les personnes séropositives ne souffrant d'aucune autre MST et suivant un traitement antirétroviral efficace ne transmettent pas le VIH par voie sexuelle. *Bulletin des médecins suisses* 2008;89: 165-169.

Verweel G, Sharland M, Lyall H, et al. Nevirapine use in HIV-1 infected children. *AIDS* 2003; 17: 1639-47.

Wang S, Jiang P, Peng G. HBV transmission from father to foetus and HBV DNA in tissues outside the liver. *Zhonghua Gan Zang Bing Za Zhi* 1999;7:203-206.

Wohl DA, McComsey G, Tebas P, Brown TT, Glesby MJ, Reeds D, et al. Current concepts in the diagnosis and management of metabolic complications of HIV infection and its therapy. *Clin Infect Dis* 2006;43:645-653.

Wright TC, Schiffman M, Solomon D, et al. Interim Guidance for the Use of Human Papillomavirus DNA Testing as an Adjunct to Cervical Cytology for Screening. *Obstetrics and Gynecology* 2004; 103:304-309.

Xu C, Politch JA, mayer KH, Anderson DJ. Human immunodeficiency virus type-I episomal cDNA in semen. *AIDS Research and Therapy* 2005: 2:9-15